



P75. AVALIAÇÃO DIAGNOSTICA DE MASSA PELVICA.

de Sousa J D (BR) [1], Machado Y d (BR) [2], Simoes M R (BR) [3], Baratella B M (BR) [4]

PURPOSE: To diagnose pelvic mass in women at menacme.

METHODS: Case report.

PATIENT: D.A.M, aged 31, from Porto Velho, Rondônia. It presents complaint of pain region of left iliac fossa approximately 1 year. Transvaginal ultrasonography revealed left adnexal tumor with negative tumor markers.

INTERVENTION: During his stay at the Hospital of Base Ary Pinheiro, a physical examination with a palpable mass was present in the left iliac fossa, irregularly, with a mean and mobile volume, with no signs of ascites and painful palpation, gynecological examination without alterations. Who underwent further examinations, found that she presented a case of pelvic kidney in topography of the left attachment.

RESULTS: New ultrasonography and tumor markers were performed, which maintained the hypothesis of pelvic kidney, submitted to excretory urography confirming hypothesis of pelvic kidney and negative markers.

CONCLUSION: The importance of the correlation of physical examination and complementary tests, for a faithful diagnosis.

[1] Médica Residente do Hospital de Base Dr. Ary Pinheiro, [2] Secretaria de Estado da Saúde de Rondônia - SESAU RO, [3] Coordenadora Programa de Residência Médica em Ginecologia e Obstetrícia do Hospital de Base Dr. Ary Pinheiro, [4] Médica Residente Hospital de Base Dr Ary Pinheiro